

## Table Notes

### Heart Attack – Angioplasty/Stent

**Total Number of Cases** includes all inpatient hospitalizations, after exclusions, for patients 18 years and older who underwent an angioplasty/stent procedure for heart attack. Patients who were transferred to another acute care hospital or left against medical advice were excluded.

**Mortality** represents patients who died during the hospital stay.

**Readmission** represents patients who were readmitted to a Pennsylvania general or specialty general acute care hospital within 30 days of the discharge date of the original hospitalization. Out-of-state residents were excluded because readmission data was not available for patients readmitted to a non-Pennsylvania hospital. Planned readmissions were not counted.

**Average Hospital Charge** represents the entire length of stay and is trimmed and case-mix adjusted. Professional fees were not included. In almost all cases, hospitals typically receive actual payments from private insurers or government payers that are considerably less than the listed charge.

*These procedures are used to open narrowed or blocked coronary (heart) arteries to restore blood flow to the heart muscle by inserting a catheter (thin tube) through the skin of the upper thigh or arm. An angioplasty is performed by inflating a small balloon at the end of the catheter to widen the narrowed or blocked area in the artery. Angioplasty procedures are often combined with the placement of a stent (permanent metallic mesh tube) to prevent the artery from narrowing again. Only patients who had a myocardial infarction (heart attack) are included.*

#### Understanding the Symbols

The symbols displayed in this report represent a comparison of a hospital's actual mortality (or readmission) rate to what is expected, after accounting for patient risk.

- **Rate was significantly lower than expected.** Fewer patients died (or were readmitted) than could be attributed to patient risk and random variation.
- ◉ **Rate was not significantly different than expected.** The number of patients who died (or were readmitted) was within the range anticipated based on patient risk and random variation.
- **Rate was significantly higher than expected.** More patients died (or were readmitted) than could be attributed to patient risk and random variation.

See **About the Report** section or **Technical Notes** for further details.