
Transcatheter Aortic Valve Replacement (TAVR)



This report examines the **8,618** hospitalizations for transcatheter aortic valve replacement (TAVR) procedures performed in Pennsylvania acute care hospitals during Calendar Years (CY) 2019 and 2020.

Of these hospitalizations, **8,412** cases, after exclusions, were examined in the Hospital Results section of this report in determining statewide rates of readmission and postoperative length of stay.



Mortality

- The in-hospital mortality rate was **0.7%**.
- Mortality ratings are not reported for hospitals and are not displayed in the Hospital Results section of the report.

Readmission

- **4.9%** of patients were readmitted to an acute care hospital within 7 days, **10.5%** were readmitted within 30 days, and **17.7%** were readmitted within 90 days.
- Hispanic patients had the highest 30-day readmission rate at **16.0%**. Black (non-Hispanic) and White (non-Hispanic) patients had 30-day readmission rates of **12.3%** and **10.6%**, respectively.

Pennsylvania Health Care Cost Containment Council (PHC4) Cardiac Procedures Report • January 2019 through December 2020 Data Transcatheter Aortic Valve Replacement (TAVR) • Page 1

Postoperative Length of Stay

- In CY 2019-2020, patients stayed in the hospital an average of **2.3** days following a TAVR procedure.
- 4.9% of patients experienced an extended postoperative length of stay; that is, after accounting for patient risk, they stayed in the hospital longer than expected. The average postoperative stay for these patients was 10.0 days.

Medicare Payments

- The average Medicare fee-for-service payment for hospitalizations in CY 2019-2020 was \$45,184.
- The average Medicare fee-for-service payment for hospitalizations with an extended postoperative length of stay was \$58,624.

Based on the Centers for Medicare and Medicaid Services CY 2019-2020 Medicare fee-for-service payment data.

TAVR Hospitalization Rates per 10,000 Pennsylvania Residents (CY 2019-2020) Statewide Rate: 7.6

Age	Female/Male	Race/Ethnicity
Age 18-640.5 Age 65-7412.4 Age 75-8447.8 Age 85 and older65.5	Female 6.7 Male 8.6	White (non-Hispanic)9.0 Black (non-Hispanic)2.1 Hispanic1.2
Poverty	Rural/Urban	Region

Rate calculations: Hospitalization rates above were based on Pennsylvania residents (7,715 hospital stays or 89.5% of all 8,618 hospitalizations for TAVR) using PHC4 discharge data and 2020 US Census Bureau population figures, with the exception of rates by poverty level, which were based on 2019 US Census Bureau estimates.

High Poverty rates included hospitalizations for residents living in areas where 20% or more of the population lives in poverty.

Rural County and *Urban County* rates were based on the designation of the county of residence as defined by The Center for Rural Pennsylvania. Urban counties are italicized in the following list of counties by region. Regional rates were adjusted for population differences in age and sex.

- Western PA includes the following counties: Allegheny, Armstrong, Beaver, Bedford, Blair, Butler, Cambria, Cameron, Clarion, Clearfield, Crawford, Elk, Erie, Fayette, Forest, Greene, Indiana, Jefferson, Lawrence, McKean, Mercer, Potter, Somerset, Venango, Warren, Washington, and Westmoreland.
- Central & Northeastern PA includes the following counties: Adams, Bradford, Centre, Clinton, Columbia, *Cumberland, Dauphin*, Franklin, Fulton, Huntingdon, Juniata, *Lackawanna, Lancaster, Lebanon, Luzerne*, Lycoming, Mifflin, Monroe, Montour, Northumberland, Perry, Pike, Snyder, Sullivan, Susquehanna, Tioga, Union, Wayne, Wyoming, and *York*.
- Southeastern PA includes the following counties: Berks, Bucks, Carbon, Chester, Delaware, Lehigh, Montgomery, Northampton, Philadelphia, and Schuylkill.

For County Rates, see "Maps – Rates by County"

Pennsylvania Health Care Cost Containment Council (PHC4) Cardiac Procedures Report • January 2019 through December 2020 Data Transcatheter Aortic Valve Replacement (TAVR) • Page 2